BOVINE ACCELLYTE II

SOURCE OF ENERGY, ELECTROLYTES AND OSMOLYTES FOR HYDRATION SUPPORT OF YOUNG, GROWING AND ADULT CATTLE

For optimal absorption of electrolytes and water



Unique formulation designed for adult animals and young stock to promote electrolyte balance for optimal hydration

 Helps to promote feed and water intake which helps restore electrolyte functions and water retention to prevent losses due to dehydration

Formulated with proper balance of electrolyte combination

- Useful during periods of stress such as:
 - Heat stress
 - Freshening
 - Feed changes
 - Illness/sick pen
 - Transportation
- Helps to maintain production and reduce panting and other heat stress related symptoms

Allows for multiple applications

- Can be dry-mixed into total mixed ration (TMR) for group or herd application
- Mixes easily with water and can be added to stock tanks or individual water feedings

Flexibility of use

 For heat stress, can be fed continually throughout the summer months or can be added to diets two days prior to any stressful event and continued through the stressful period

AVAILABLE PACKAGE SIZES

OVINE ACCELLYTE II

10 lb. pail

- Contains 160 1 oz. feedings
- Case = 12 10 lb. pails

25 lb. bags

- Contains 400 1 oz. feedings
- Pallet = 80 25 lb. bags

STORAGE

• Store in a cool, dry place. Keep tightly closed.

DIRECTIONS FOR USE

Adult Cattle Application

Top-dress or mix into TMR or grain portion of the diet or add to water supply at the rate of:

- Feed 1-2 ounces per head per day. Optimally, in the first TMR feeding of each day when animals
 are the coolest and at their strongest feed intakes.
- 2-3 ounces per head per day for three to seven days prior to shipping or after receiving cattle.

Show Ring Cattle Application

Top-dress or mix into TMR or grain portion of the diet or add to water supply at the rate of:

• 1-2 ounces per head per day for one week before transport, continuing throughout the show and two days after returning.

Young and Growing Cattle Application

Top-dress or mix into TMR or grain portion of the diet or add to water supply at the rate of:

 ¼-½ ounce per head per day with daily feed or water supply during heat stress conditions or anytime dehydration is at risk.

USAGE GUIDE

The chart to the right illustrates product usage based on the mature cattle application rate of 1 ounce per head per day.

Feeding Period	1 Day	2 Weeks	1 Month
50 Cows/Adult Animals	3.1 lbs.	43.4 lbs.	93 lbs.
75 Cows/Adult Animals	4.7 lbs.	65.6 lbs.	141 lbs.
100 Cows/Adult Animals	6.25 lbs.	87.5 lbs.	187.5 lbs.
200 Cows/Adult Animals	12.5 lbs.	175 lbs.	375 lbs.
400 Cows/Adult Animals	25 lbs.	350 lbs.	750 lbs.

