

# BUILD A BETTER CALF<sup>®</sup> ACCEL MILK FORTIFIER *MEDICATED*

## CONCENTRATED VITAMIN AND MINERAL SUPPLEMENT FOR YOUNG CATTLE

Contains a source of live (viable) naturally occurring microorganisms for control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in young calves



### AVAILABLE PACKAGE SIZES

12 lb. bag

25 lb. bag

- Pallet = 80 - 25 lb. bags

### STORAGE

- Store in a cool, dry place. Keep tightly closed.

### RESEARCH-SUPPORTED FEATURES AND BENEFITS

Contains National Research Council recommended levels of bioavailable vitamins and trace minerals

- Supplements pasteurized waste milk (PWM) deficiencies of essential nutrients which support immune development, calf health and growth

Formulated with 1 gram/head/day dosage of a probiotic source, Tri-Mic WD

- Provides beneficial microflora for a competitive exclusion of pathogens, reduction of scours, improved feed intake and utilization, and weight gain

Medicated with Bovatec<sup>®</sup>

- For the control of Coccidiosis

Low and economical 9 gram/head/day feeding rate

- Easily adaptable to PWM feeding programs

Contains ClariFly<sup>®</sup>

- Feed-through larvicide fly control
- Control house flies, face flies, stable flies and horn flies

### DIRECTIONS FOR USE

Mix 4.5 grams of Accel Milk Fortifier into 2 quarts of pasteurized waste milk and mix thoroughly (see larger batch mixing directions in the chart below). Feed twice daily. Feed immediately after mixing. Each quart of pasteurized waste milk will then contain 11.36 mg of Lasalocid. Feed to provide each calf 1 mg. of Lasalocid per 2.2 lb. (1 kg) of body weight per day. Feeding 9 g of Accel Milk Fortifier powder daily will medicate 100 lbs. of body weight.

### BATCH MIXING

Number of Calves on Milk	Grams per Feeding
1	4.5 grams
20	90 grams (3 oz.)
40	180 grams (6 oz.)
60	270 grams (9 oz.)
80	360 grams (12 oz.)
100	450 grams (15 oz.)
160	720 grams (24 oz.)
200	900 grams (30 oz.)

Above chart based on 100 lb. calf

### CAUTION

Follow label directions. The addition to feed higher levels of this premix containing selenium is not permitted. The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. Type B feeds (cattle and sheep), mix thoroughly prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration that could be fatal to cattle and sheep. When preparing large batches, keep the mixture agitated while dispensing into individual containers to prevent settling of Lasalocid.

### WARNING

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.



**YOUR SUCCESS** *Our Passion.*

(614) 873-4683 ♦ [www.selectsires.com](http://www.selectsires.com)